

Periodical volume

Mathematische Zeitschrift - 221

in: Periodical

691 page(s)

Nutzungsbedingungen

DigiZeitschriften e.V. gewährt ein nicht exklusives, nicht übertragbares, persönliches und beschränktes Recht auf Nutzung dieses Dokuments. Dieses Dokument ist ausschließlich für den persönlichen, nicht kommerziellen Gebrauch bestimmt. Das Copyright bleibt bei den Herausgebern oder sonstigen Rechteinhabern. Als Nutzer sind Sie nicht dazu berechtigt, eine Lizenz zu übertragen, zu transferieren oder an Dritte weiter zu geben.

Die Nutzung stellt keine Übertragung des Eigentumsrechts an diesem Dokument dar und gilt vorbehaltlich der folgenden Einschränkungen:

Sie müssen auf sämtlichen Kopien dieses Dokuments alle Urheberrechtshinweise und sonstigen Hinweise auf gesetzlichen Schutz beibehalten; und Sie dürfen dieses Dokument nicht in irgend einer Weise abändern, noch dürfen Sie dieses Dokument für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, aufführen, vertreiben oder anderweitig nutzen; es sei denn, es liegt Ihnen eine schriftliche Genehmigung von DigiZeitschriften e.V. und vom Herausgeber oder sonstigen Rechteinhaber vor.

Mit dem Gebrauch von DigiZeitschriften e.V. und der Verwendung dieses Dokuments erkennen Sie die Nutzungsbedingungen an.

Terms of use

DigiZeitschriften e.V. grants the non-exclusive, non-transferable, personal and restricted right of using this document. This document is intended for the personal, non-commercial use. The copyright belongs to the publisher or to other copyright holders. You do not have the right to transfer a licence or to give it to a third party.

Use does not represent a transfer of the copyright of this document, and the following restrictions apply:

You must abide by all notices of copyright or other legal protection for all copies taken from this document; and You may not change this document in any way, nor may you duplicate, exhibit, display, distribute or use this document for public or commercial reasons unless you have the written permission of DigiZeitschriften e.V. and the publisher or other copyright holders.

By using DigiZeitschriften e.V. and this document you agree to the conditions of use.

Kontakt / Contact

DigiZeitschriften e.V.

Papendiek 14

37073 Goettingen

Email: info@digizeitschriften.de

Theta functions and cycles on some abelian fourfolds

Bert van Geemen

Department of Mathematics, Ruu, P.O. Box 80.010, 3508 TA Utrecht, The Netherlands

Received 10 March 1994; in final form 15 August 1994

1 Introduction

1.1 Let X be an abelian variety over \mathbb{C} and let K be an imaginary quadratic field with

$$K \subset \text{End}(X) \otimes \mathbb{Q}.$$

The action t of K on T_0X , the tangent space at the origin of X , can be diagonalized thus, w.r.t. a suitable basis of T_0X :

$$t(k) = \text{diag}(\sigma(k), \dots, \sigma(k), \bar{\sigma}(k), \dots, \bar{\sigma}(k)), \quad (k \in K)$$

with $\sigma : K \hookrightarrow \mathbb{C}$ an embedding of K . We say X is of type (p, q) if there are p entries $\sigma(k)$ and q entries $\bar{\sigma}(k)$.

In case X has type (p, p) , Weil [W] constructed a two dimensional subspace $W \subset B^p(X) := H^{2p}(X, \mathbb{Q}) \cap H^{p,p}(X)$. For general X , $\dim X > 2$, of type (p, p) one has in fact:

$$\dim NS(X)_{\mathbb{Q}} = \dim B^1(X) = 1, \quad B^p(X) = D^p(X) \oplus W,$$

here $D^p(X) := \text{Im}(B^1(X)^p \rightarrow B^p(X))$, $(\omega_1, \dots, \omega_p) \mapsto \omega_1 \wedge \dots \wedge \omega_p$. In particular such an X has Hodge classes which are not obtained as intersections (cup products) of divisors classes. In [vG2] we gave an elementary introduction to these abelian varieties.

1.2 For certain abelian varieties X of Weil-type with field $K = \mathbb{Q}(i)$, $\mathbb{Q}(\omega)$, $\omega^3 = -1$, C. Schoen [S] gave a construction for algebraic cycles on X whose classes span W .

In this paper we use theta functions to construct algebraic cycles on certain abelian 4-folds of Weil type. In fact we show that in the case under consideration the abelian varieties allow a rational map onto a smooth quadric \bar{Q} in \mathbb{P}^5 , the classes of the inverse images of the rulings of \bar{Q} give classes which do not lie in D^2 .

Theorem 3.7 *Let X be an abelian 4-fold of Weil-type $(2, 2)$, with field $\mathbb{Q}(i)$ and $\det H = 1$.*

Then the space W of Weil–Hodge cycles is spanned by classes of algebraic cycles.

1.3 We refer to 2.1 and [vG2] for the explanation of ‘ $\det H$ ’. It turns out, see 5.2, that the abelian varieties we consider are among those studied by Schoen [S], however, our method is different. We don’t know how the cycles obtained by us and Schoen are related.

2 The abelian varieties

2.1 The abelian varieties of Weil-type we consider were introduced in [vG1], 10.6, 10.7. Let \mathbb{S}_{2n} be the Siegel upper half space of $(2n) \times (2n)$ complex, symmetric matrices whose imaginary part defines a positive definite quadratic form on \mathbb{R}^{2n} and let

$$\mathbb{H}_{2n} := \left\{ \tau = \begin{pmatrix} \tau_1 & \tau_{12} \\ -\tau_{12} & \tau_1 \end{pmatrix} \in \mathbb{S}_{2n} : \tau_1 \in \mathbb{S}_n, {}^t\tau_{12} = -\tau_{12} \right\}$$

and

$$\Omega_\tau := \begin{pmatrix} I \\ \tau \end{pmatrix}.$$

We will write vectors as row vectors. The abelian varieties are then defined by: $X = X_\tau := \mathbb{C}^{2n}/(\mathbb{Z}^{4n}\Omega_\tau)$ with $\tau \in \mathbb{H}_{2n}$. The equality:

$$M\Omega_\tau = \Omega_\tau A, \quad \text{with } M := \begin{pmatrix} 0 & I \\ -I & 0 \\ & 0 & I \\ & -I & 0 \end{pmatrix}, \quad A := \begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix},$$

shows that X_τ has an automorphism ϕ of order four with:

$$\phi : X_\tau \rightarrow X_\tau, \quad \phi_* = M \in \text{End}(H_1(X_\tau, \mathbb{Z})), \quad d\phi = A \in \text{End}(T_0 X_\tau).$$

Note that the eigenvalues of $d\phi$ are i and $-i$, both with multiplicity n , thus the X_τ are of Weil-type (n, n) .

The principal polarization $E : \wedge^2 H_1(X, \mathbb{Z}) \rightarrow \mathbb{Z}$ on X_τ is given by an alternating $4n \times 4n$ matrix of the same shape as A , but with four $2n \times 2n$ blocks rather than $n \times n$ blocks. A simple computation verifies that ϕ_* preserves the polarization:

$$E(\phi_* x, \phi_* y) = E(x, y) \quad (\forall x, y \in H_1(X, \mathbb{Z})).$$

The space $H_1(X, \mathbb{Q})$ has the structure of a $\mathbb{Q}(i)$ -vector space via the action of ϕ_* . Define:

$$H : H_1(X, \mathbb{Q}) \times H_1(X, \mathbb{Q}) \rightarrow \mathbb{Q}(i); \quad H(x, y) := E(x, \phi_* y) + iE(x, y).$$

Then H is a Hermitian form ($H(x, y) = \overline{H(y, x)}$ and $H(x, \phi_* y) = iH(x, y)$) and its determinant (modulo norms of elements of \mathbb{Q}^*) is an isogeny invariant. An easy computation shows that

$$\det H = 1.$$

2.2 We recall the definition of theta functions with characteristics $m, m' \in \mathbb{R}^g$ (cf. [I], p. 49):

$$\theta_{m, m'}(\tau, z) = \sum_{k \in \mathbb{Z}^g} \exp(\pi i[(k + m)\tau'(k + m) + 2(k + m)'(z + m')]),$$

here $\tau \in \mathbb{S}_g$, $z \in \mathbb{C}^g$. The zero locus of the function $\theta_{0,0}(\tau, z)$ (Riemann's theta function on X_τ) in \mathbb{C}^g defines a symmetric divisor $\Theta_\tau \subset X_\tau$. The pair $(X_\tau, \mathcal{O}(\Theta_\tau))$ is a principally polarized abelian variety. Thus the Riemann theta function is the pull-back of a global section of $\mathcal{O}(\Theta_\tau)$ to \mathbb{C}^g . If $X_\tau \cong X_{\tau'}$ one has $\mathcal{O}(2\Theta_\tau) \cong \mathcal{O}(2\Theta_{\tau'})$, thus this line bundle is canonically defined (intrinsically: the unique totally symmetric line bundle algebraically equivalent to twice the principal polarization).

We define theta functions with half integral characteristics:

$$\theta \begin{bmatrix} m \\ m' \end{bmatrix}(\tau, z) := \theta_{m/2, m'/2}(\tau, z) \quad \text{for } m, m' \in \{0, 1\}^g.$$

Let α the point of order two in X_τ defined by $(m + \tau m')/2 \in \mathbb{C}^g$ and let $(\Theta_\tau)_\alpha$ be the translate of Θ_τ by α . Then the function $\theta \begin{bmatrix} m \\ m' \end{bmatrix}$ is a basis of $H^0(X_\tau, \mathcal{O}((\Theta_\tau)_\alpha))$. The 2^g functions:

$$\theta \begin{bmatrix} m \\ 0 \end{bmatrix}(2\tau, 2z), \quad m \in \{0, 1\}^g$$

are a basis of $H^0(X_\tau, \mathcal{O}(2\Theta_\tau))$.

From the definition of the theta function one easily proves the following lemma, which gives the action of ϕ^* on global sections of certain line bundles on X_τ and which gives a formula for the maps

$$(1 - \phi)^* : H^0(X, \mathcal{O}(\Theta_\alpha)) \rightarrow H^0(X, \mathcal{O}(2\Theta)),$$

for the translates Θ_α of Θ with $\alpha \in \ker(1 - \phi)$.

2.3 Lemma. For $\varepsilon_i, \varepsilon'_i \in \{0, 1\}^n$, $z \in \mathbb{C}^{2n}$, $\tau \in \mathbb{H}_{2n}$ and $N \in \mathbb{Z}_{\geq 0}$ we have:

1.

$$\theta \begin{bmatrix} \varepsilon_1 & \varepsilon_2 \\ \varepsilon'_1 & \varepsilon'_2 \end{bmatrix}(N\tau, zA) = (-1)^{\varepsilon'_1 \varepsilon'_1} \theta \begin{bmatrix} \varepsilon_2 & \varepsilon_1 \\ \varepsilon'_2 & \varepsilon'_1 \end{bmatrix}(N\tau, z).$$

2.

$$\theta \begin{bmatrix} \varepsilon & \varepsilon \\ \varepsilon' & \varepsilon' \end{bmatrix}(N\tau, z(I - A)) = \sum_{\rho \in \{0, 1\}^n} (-1)^{(\rho + \varepsilon)' \varepsilon'} \theta \begin{bmatrix} \rho & \rho + \varepsilon \\ 0 & 0 \end{bmatrix}(2N\tau, 2z).$$

Proof. First we prove the second statement. For $\tau \in \mathbb{H}_{2n}$ (and ${}^t A$ the transposed of A):

$$A^2 = -I, \quad {}^t A = -A, \quad A\tau' A = \tau, \quad A\tau + \tau' A = 0.$$

Note that these relations remain valid upon replacing τ by $N\tau$, we will suppress N in the rest of the proof.

We compute $\theta_{(m_1, m_2), (m'_1, m'_2)}(\tau, z(I - A))$. Using the relations, we can write:

$$\tau = (I + A)(1/2)\tau'(I + A), \quad I = (1/2)(I + A)(I - A) .$$

and we get:

$$(k + m)\tau'(k + m) = (k + m)(I + A)(1/2)\tau'(I + A)'(k + m) ,$$

$$(k + m)'(z(I - A) + m'I) = (k + m)(I + A)'(z + (m'/2)(I + A)) .$$

Writing $k = (k_1, k_2)$, $m = (m_1, m_2)$, $m' = (m'_1, m'_2) \in \mathbb{Z}^n \times \mathbb{Z}^n$ we get:

$$\begin{aligned} (k + m)(I + A) &= (k_1 - k_2 + m_1 - m_2, k_1 + k_2 + m_1 + m_2) \\ &= (2l_1 + \rho + m_1 - m_2, 2l_2 + \rho + m_1 + m_2) , \end{aligned}$$

with $\rho \in \{0, 1\}^n$ defined by $k_1 \pm k_2 \in \rho + 2\mathbb{Z}^n$ and $l_1 := (k_1 - k_2 - \rho)/2$, $l_2 := (k_1 + k_2 - \rho)/2 \in \mathbb{Z}^n$. Then:

$$(k + m)\tau'(k + m) = (l_1 + (\rho + m_1 - m_2)/2, l_2 + (\rho + m_1 + m_2)/2)(2\tau)'(l_1 + \dots)$$

and in the second term we get:

$$(z + (m'/2)(I + A)) = (2z + (m'_1 - m'_2, m'_1 + m'_2)/2) ,$$

thus by summation of the terms over $\rho \in \{0, 1\}^n$, $l_1, l_2 \in \mathbb{Z}^n$ we have:

$$\begin{aligned} &\theta_{(m_1, m_2), (m'_1, m'_2)}(\tau, z(I - A)) \\ &= \sum_{\rho \in \{0, 1\}^n} \theta_{(m_1 - m_2 + \rho, m_1 + m_2 + \rho)/2, (m'_1 - m'_2, m'_1 + m'_2)/2}(2\tau, 2z) . \end{aligned}$$

The isogeny formula follows from this, but since we want (00) rather than $(0\epsilon')$ in the second row of the characteristic, we get the sign (cf. [I], $\theta.2$, p.49).

For the first formula, we observe that

$$(k + m)'(zA + m') = (k + m)'A'(z + m'A^{-1}),$$

$$(k + m)\tau'(k + m) = (k + m)'A\tau A'(k + m) ,$$

and that

$$(k + m)'A = (k_2 + m_2, -k_1 - m_1), \quad m'A^{-1} = (k_2 + m_2, -k_1 - m_1) .$$

Thus we have:

$$\theta_{(m_1, m_2), (m'_1, m'_2)}(\tau, zA) = \theta_{(m_2, -m_1), (m'_2, -m'_1)}(\tau, z) .$$

Since we want characteristics in $\{0, 1\}^{2n}$, we get the sign in the formula. \square

2.4 Corollary. *The automorphism ϕ of order two induces an automorphism ϕ^* on $H^0(X_\tau, \mathcal{O}(2\Theta_\tau))$ of order two (in fact, the corresponding theta functions are even). Let $H^0(X_\tau, \mathcal{O}(2\Theta_\tau))_\pm$ be the \pm eigenspace of ϕ^* , then*

$$\begin{aligned}\dim H^0(X_\tau, \mathcal{O}(2\Theta_\tau))_+ &= 2^{n-1}(2^n + 1), \\ \dim H^0(X_\tau, \mathcal{O}(2\Theta_\tau))_- &= 2^{n-1}(2^n - 1).\end{aligned}$$

Bases of $\dim H^0(X_\tau, \mathcal{O}(2\Theta_\tau))_\pm$ are given by the non-zero

$$\theta \begin{bmatrix} \varepsilon_1 & \varepsilon_2 \\ 0 & 0 \end{bmatrix} (2\tau, 2z) \pm \theta \begin{bmatrix} \varepsilon_2 & \varepsilon_1 \\ 0 & 0 \end{bmatrix} (2\tau, 2z)$$

with $\varepsilon_1, \varepsilon_2 \in \{0, 1\}^n$.

2.5 Remark. The isogeny formula 2.3.2 implies that each of the functions $\theta \begin{bmatrix} \sigma \\ 0 \end{bmatrix} (2\tau, 2z)$, ($\sigma \in \{0, 1\}^{2n}$) can be expressed as a linear combination of the 2^{2n} functions $\theta \begin{bmatrix} \varepsilon & \varepsilon' \\ \varepsilon & \varepsilon' \end{bmatrix} (\tau, z(I - A))$, ($\varepsilon, \varepsilon' \in \{0, 1\}^n$). Thus these functions also provide a basis of $H^0(X_\tau, \mathcal{O}(2\Theta_\tau))$.

3 The cycles

3.1 In this section, $X = X_\tau$ with $\tau \in \mathbb{H}_{2n}$. Let $N = 2^{2n} - 1$. We study the geometry of the Kummer map:

$$\begin{aligned}K : X_\tau &\rightarrow \mathbb{P}^N = \mathbb{P}H^0(X, \mathcal{O}(2\Theta)), \\ z &\mapsto (\cdots : \theta \begin{bmatrix} \sigma \\ 0 \end{bmatrix} (2\tau, 2z) : \cdots) \quad (\sigma \in \{0, 1\}^{2n}).\end{aligned}$$

In case X is indecomposable, the map K embeds $X/\pm 1$ in \mathbb{P}^N , if the ppav is a product of ppav's $X = X_1 \times \cdots \times X_k$ then $K(X) \cong K(X_1) \times \cdots \times K(X_k)$.

3.2 The morphism K is equivariant for the action of the automorphism of order 4, which acts on \mathbb{P}^N by (see 2.3.1):

$$(\cdots : X_{\sigma_1 \sigma_2} : \cdots) \mapsto (\cdots : X_{\sigma_2 \sigma_1} : \cdots) \quad (\sigma_1, \sigma_2 \in \{0, 1\}^n).$$

The two eigenspaces of this involution we denote by (cf. 2.4)

$$\mathbb{P}_\pm := \mathbb{P}H^0(X_\tau, \mathcal{O}(2\Theta_\tau))_\pm$$

Let $\Pi_- : \mathbb{P}^N \rightarrow \mathbb{P}_-$ be the projection on to \mathbb{P}_- from \mathbb{P}_+ , and let:

$$K_- := \Pi_- \circ K : X \rightarrow \mathbb{P}_-,$$

be the composition of K with the projection Π_- .

3.3 Lemma. *For an indecomposable ppav the base locus B of the rational map $K_- : X \rightarrow \mathbb{P}_-$ consists of a set of 2^{2n} two-torsion points, in fact:*

$$B = \ker(\phi - 1) = \{x \in X : \phi(x) = x\} \cong (\mathbb{Z}/2\mathbb{Z})^{2n}.$$

Moreover, B is a maximal isotropic subgroup of $X[2]$ and $X/B \cong X$.

Proof. Since K is a morphism, $K(B)$ is base locus of Π_- , so $K(B) = K(X) \cap \mathbb{P}_+$. As K is equivariant for ϕ and \mathbb{P}_+ is an eigenspace, B is contained in the set of points $x \in X$ with $\phi(x) = \pm x$, that is $(\phi \pm 1)x = 0$. Note $\phi \circ (\phi + 1) = \phi - 1$, so $\ker(\phi + 1) = \ker(\phi - 1)$; since $(\phi - 1)^2 = -2\phi$ we get $B \subset \ker(\phi - 1) \cong (\mathbb{Z}/2\mathbb{Z})^{2n}$.

To see that all points in $\ker(\phi - 1)$ are mapped to \mathbb{P}_+ (and not to \mathbb{P}_-), note first of all that $0 \in X$ is mapped to \mathbb{P}_+ since 2.3.1 (with $N = 2$ there) implies that $\theta \begin{bmatrix} \sigma_1 & \sigma_2 \\ 0 & 0 \end{bmatrix} (2\tau, 0) = \theta \begin{bmatrix} \sigma_2 & \sigma_1 \\ 0 & 0 \end{bmatrix} (2\tau, 0)$.

The map K is equivariant for translations by points of order two (they act on \mathbb{P}^N as the Heisenberg group), thus each $a \in X[2]$ gives a projective transformation $U(a)$ satisfying

$$K(x + a) = U(a)K(x) \quad (a \in X[2]).$$

Let $a \in \ker(\phi - 1)$. Then translation by a and ϕ commute ($\phi(x + a) = \phi(x) + \phi(a) = \phi(x) + a$ for all $x \in X$). Therefore $U(a)$ maps the eigenspaces of ϕ into themselves (it cannot interchange them since the dimensions are different). Thus $K(a) = U(a)K(0)$ also lies in \mathbb{P}_+ and $B = \ker(\phi - 1)$.

That $X/B \cong X$ is trivial since $\phi - 1 \in \text{End}(X)$. Note that $B = \ker(\phi - 1) = \text{im}(\phi - 1)$, $\phi_*^2 = -1$ and $E(\phi_*x, \phi_*y) = E(x, y)$ give:

$$E((\phi_* - 1)x, (\phi_* - 1)y) = 2E(x, y) + E(\phi_*^2x, \phi_*y) - E(\phi_*x, y) = 2E(x, y).$$

The polarization E defines the Weil-pairing and this formula shows that the Weil-pairing is trivial on B . \square

3.4 Note that the proof shows that $K(X) \cap \mathbb{P}_-$ is empty, so the projection of $K(X)$ to \mathbb{P}_+ is a morphism.

3.5 We recall the basic facts on the quadrics on \mathbb{P}^N . A basis for the vector space of these quadrics is provided by the

$$Q \begin{bmatrix} \varepsilon \\ \varepsilon' \end{bmatrix} := \sum_{\sigma} (-1)^{\sigma' \varepsilon'} X_{\sigma} X_{\sigma + \varepsilon}, \quad (\sigma, \varepsilon, \varepsilon' \in (\mathbb{Z}/2\mathbb{Z})^{2n}, \varepsilon' \varepsilon' = 0),$$

which are indexed by the $2^{2n-1}(2^{2n} + 1)$ even theta characteristics (each $Q_{\begin{bmatrix} \varepsilon \\ \varepsilon' \end{bmatrix}}$ is an eigenvector for the action of the Heisenberg group).

The pull-back of a quadric Q along K gives a section of $\mathcal{O}(4\Theta)$ and, since the $\theta \begin{bmatrix} \sigma \\ 0 \end{bmatrix} (2\tau, 2z)$ are even functions in z , this section will also be given by an even theta function. A basis of $H^0(X_{\tau}, \mathcal{O}(4\Theta))_+$, the space of these even theta functions, is given by the $\theta \begin{bmatrix} \varepsilon \\ \varepsilon' \end{bmatrix} (\tau, 2z)$ with $\varepsilon' \varepsilon' = 0$. The following theta relation ([I], IV.1) expresses the pull-back of the $Q_{\begin{bmatrix} \varepsilon \\ \varepsilon' \end{bmatrix}}$'s in this basis:

$$\begin{aligned} K^* Q_{\begin{bmatrix} \varepsilon \\ \varepsilon' \end{bmatrix}} &:= \sum_{\sigma} (-1)^{\sigma' \varepsilon'} \theta \begin{bmatrix} \sigma \\ 0 \end{bmatrix} (2\tau, 2z) \theta \begin{bmatrix} \sigma + \varepsilon \\ 0 \end{bmatrix} (2\tau, 2z) \\ &= \theta \begin{bmatrix} \varepsilon \\ \varepsilon' \end{bmatrix} (\tau, 0) \theta \begin{bmatrix} \varepsilon \\ \varepsilon' \end{bmatrix} (\tau, 2z). \end{aligned}$$

In particular, $K(X)$ lies in $Q_{[\frac{\varepsilon}{\varepsilon'}]}$ (we denote the quadric in \mathbb{P}^N and its defining equation by the same symbol) iff the corresponding theta constant vanishes, i.e. $\theta[\frac{\varepsilon}{\varepsilon'}](\tau, 0) = 0$.

From Lemma 2.3.1 we have that on X_τ the $2^{n-1}(2^n - 1)$ even theta constants $[\frac{\varepsilon}{\varepsilon'} \frac{\varepsilon'}{\varepsilon'}]$ with $\varepsilon' \varepsilon' \equiv 1 \pmod{2}$ vanish.

In the case $n = 2$ the 6 even theta constants with the following characteristics vanish:

$$\begin{bmatrix} 1010 \\ 1010 \end{bmatrix}, \begin{bmatrix} 1010 \\ 1111 \end{bmatrix}, \begin{bmatrix} 1111 \\ 1010 \end{bmatrix}, \begin{bmatrix} 1111 \\ 0101 \end{bmatrix}, \begin{bmatrix} 0101 \\ 0101 \end{bmatrix}, \begin{bmatrix} 0101 \\ 1111 \end{bmatrix}.$$

Therefore $K(X)$ lies in 6 independent quadrics. We will show, by explicit computation, that there exists a (unique) quadric Q which is linear combination of these 6 quadrics and which is the cone over a smooth quadric in \mathbb{P}^5 . This implies that the image of $K(X)$ in \mathbb{P}^5 is in this quadric.

3.6 Proposition. *Define a quadric Q in \mathbb{P}^{15} by:*

$$Q := Q \begin{bmatrix} 1010 \\ 1010 \end{bmatrix} - Q \begin{bmatrix} 1010 \\ 1111 \end{bmatrix} - Q \begin{bmatrix} 1111 \\ 1010 \end{bmatrix} + Q \begin{bmatrix} 1111 \\ 0101 \end{bmatrix} - Q \begin{bmatrix} 0101 \\ 0101 \end{bmatrix} + Q \begin{bmatrix} 0101 \\ 1111 \end{bmatrix}.$$

Then for any $\tau \in \mathbb{H}_4$, we have $K(X_\tau) \subset Q$ and Q is the cone over a smooth quadric \overline{Q} in $\mathbb{P}^5 = \mathbb{P}_-$, thus:

$$K_- : X - B \rightarrow \overline{Q} \subset \mathbb{P}^5.$$

Proof. Using the definition of $Q[\frac{\varepsilon}{\varepsilon'}]$, it is clear that $Q[\frac{1010}{1010}]$ and $Q[\frac{1010}{1111}]$ involve the same 8 monomials ($X_\sigma X_{\sigma+(1010)} = X_{0000}X_{1010}, \dots$) with coefficients ± 2 . The signs differ iff $\varepsilon' \sigma = (1010) \cdot {}^t \sigma = -1$, and thus:

$$Q \begin{bmatrix} 1010 \\ 1010 \end{bmatrix} - Q \begin{bmatrix} 1010 \\ 1111 \end{bmatrix} = 4(X_{0001}X_{1011} + X_{0100}X_{1110} - X_{0011}X_{1001} - X_{1100}X_{0110}).$$

Similarly, one finds:

$$\begin{aligned} Q \begin{bmatrix} 1111 \\ 1010 \end{bmatrix} - Q \begin{bmatrix} 1111 \\ 0101 \end{bmatrix} &= 4(X_{0001}X_{1110} + X_{0100}X_{1011} - X_{0010}X_{1101} \\ &\quad - X_{1000}X_{0111}), \\ Q \begin{bmatrix} 0101 \\ 0101 \end{bmatrix} - Q \begin{bmatrix} 0101 \\ 1111 \end{bmatrix} &= 4(X_{0010}X_{0111} + X_{1000}X_{1101} - X_{0011}X_{0110} \\ &\quad - X_{1100}X_{1001}). \end{aligned}$$

As coordinates on $\mathbb{P}^9 = \mathbb{P}_+$ we take:

$$Z_{\sigma_1 \sigma_2} := X_{\sigma_1 \sigma_2} + X_{\sigma_2 \sigma_1}$$

with the convention that in $Z_{\sigma_1\sigma_2}$ with $\sigma_1 = (a_1, b_1)$, $\sigma_2 = (a_2, b_2) \in \{0, 1\}^2$ we have that $2a_1 + b_1 \leq 2a_2 + b_2$, and on \mathbb{P}^5 we take:

$$\begin{aligned} Y_{01} &:= X_{0001} - X_{0100}, & Y_{02} &:= X_{0010} - X_{1000}, & Y_{03} &:= X_{0011} - X_{1100}, \\ Y_{12} &:= X_{0110} - X_{1001}, & Y_{13} &:= X_{0111} - X_{1101}, & Y_{23} &:= X_{1011} - X_{1110}, \end{aligned}$$

it is convenient to agree that $Y_{ab} = -Y_{ba}$ if $a > b$. Using the identity

$$2(Z_{st}Z_{uv} + Z_{ts}Z_{vu}) = (Z_{st} - Z_{ts})(Z_{uv} - Z_{vu}) + (Z_{st} + Z_{ts})(Z_{uv} + Z_{vu}),$$

with $s, t, u, v \in (\mathbb{Z}/2\mathbb{Z})^2$, we get:

$$\begin{aligned} Q \begin{bmatrix} 1010 \\ 1010 \end{bmatrix} - Q \begin{bmatrix} 1010 \\ 1111 \end{bmatrix} &= 2(Y_{01}Y_{23} + Y_{03}Y_{12} + Z_{01}Z_{23} - Z_{03}Z_{12}), \\ Q \begin{bmatrix} 1111 \\ 1010 \end{bmatrix} - Q \begin{bmatrix} 1111 \\ 0101 \end{bmatrix} &= 2(-Y_{01}Y_{23} + Y_{02}Y_{13} + Z_{01}Z_{23} - Z_{02}Z_{13}), \\ Q \begin{bmatrix} 0101 \\ 0101 \end{bmatrix} - Q \begin{bmatrix} 0101 \\ 1111 \end{bmatrix} &= 2(Y_{02}Y_{13} - Y_{03}Y_{12} + Z_{02}Z_{13} - Z_{03}Z_{12}). \end{aligned}$$

the equation for Q is then:

$$Q = 4(Y_{01}Y_{23} - Y_{02}Y_{13} + Y_{03}Y_{12}),$$

so Q is the cone over a quadric \overline{Q} in \mathbb{P}^5 . Since $K(X) \subset Q$ we get $K_-(X) \subset \overline{Q}$. \square

3.7 Theorem. *Let X be an abelian 4-fold of Weil type $(2, 2)$, field $\mathbb{Q}(i)$ and $\det H = 1$.*

Then the space W of Weil–Hodge cycles is spanned by classes of algebraic subvarieties.

Proof. Since X is isogeneous to an X_τ (cf. [vG2]), it suffices to prove the result for these varieties. We show that for a general X_τ the rational map $K_- : X_\tau \rightarrow \overline{Q}$ is dominant and that the (strict) pull-back of a general linear subspace $\mathbb{P}^2 \subset \overline{Q}$ has a cohomology class which is not in $D^2(X)$. Using the action of $\mathbb{Q}(i)^*$ by pull-back on $B^2(X) = D^2(X) \oplus W$, so $x \cdot (v, w) = ((x\bar{x})^2 v, x^4 w)$, with $W = \wedge_{\mathbb{Q}(i)}^4 H^1(X, \mathbb{Q}) \cong K$, it then follows that W is spanned by classes of cycles. Since any two points in \mathbb{H}_4 can be connected by a holomorphic curve, the closure of such a cycle in the generic fiber specializes to a cycle with a class not in $D^2(X)$ for any X_τ for $\tau \in \mathbb{H}_4$.

To show that K_- is dominant and to compute the cycle class, we specialize to the case that $X \cong Y \times Y$ with Y a general abelian surface. In terms of the period matrix for X that means we take $\tau \in \mathbb{H}_4$ with $\tau_{12} = 0$. One has:

$$H^0(X, \mathcal{O}(2\Theta_X)) \cong H^0(Y, \mathcal{O}(2\Theta_Y)) \otimes H^0(Y, \mathcal{O}(2\Theta_Y))$$

(in fact, from the definition one has: $\theta \begin{bmatrix} \sigma_1 & \sigma_2 \\ 0 & 0 \end{bmatrix} (2\tau, 2z) = \theta \begin{bmatrix} \sigma_1 \\ 0 \end{bmatrix} (2\tau_1, 2z_1) \theta \begin{bmatrix} \sigma_2 \\ 0 \end{bmatrix} (2\tau_2, 2z_2)$ with $z = (z_1, z_2) \in \mathbb{C}^2 \times \mathbb{C}^2$). The coordinate functions Y_{ij} on \mathbb{P}^5

(cf. proof of 3.6) pull back along K_- to a basis for $\wedge^2 H^0(Y, \mathcal{O}(2\Theta_Y)) \subset H^0(X, \mathcal{O}(2\Theta_X))$.

Therefore we get the following diagram:

$$\begin{array}{ccc} Y \times Y & \xrightarrow{K_-} & \overline{Q} \subset \mathbb{P}^5 \\ \downarrow K \times K & & \uparrow \cong \uparrow \cong \\ \mathbb{P}(\mathbb{C}^4) \times \mathbb{P}(\mathbb{C}^4) & \xrightarrow{P} & G(1, 3) \subset \mathbb{P}(\wedge^2 \mathbb{C}^4) \end{array}$$

(note that the horizontal maps are not everywhere defined), here $G(1, 3)$ is the Grassmanian of lines in \mathbb{P}^3 and P is the Plücker map which sends a pair of points to the line connecting them. Note that the base locus of P is exactly the diagonal of $\mathbb{P}^3 \times \mathbb{P}^3$. Since $K(Y) \cong Y/\langle \pm 1 \rangle$, the base locus of K_- is the union of the diagonal Δ of $Y \times Y$ and $\Delta^- := \{(y, -y) : y \in Y\}$. (The base locus of K_- is now larger (cf. 3.3) since $K : X \rightarrow \mathbb{P}^{15}$ factors over the group generated by $id_Y \times (-id_Y)$ and $(-id_Y) \times id_Y$.)

The equation for \overline{Q} is just the quadratic Plücker relation, it is identically zero in $\mathbb{P}^3 \times \mathbb{P}^3$. We will from now on identify $\overline{Q} = G(1, 3)$.

It is clear that K_- is dominant: to a general $l \in G(1, 3)$ corresponds a line in \mathbb{P}^3 which intersects the Kummer variety $K(Y)$ in 4 points, over each pair of these points there lie 4 elements in $Y \times Y$ mapping to l under K_- . Therefore K_- is dominant for the general X_τ with $\tau \in \mathbb{H}_4$.

A basis for $H^4(\overline{Q}, \mathbb{Q})$ is given by the classes of a $\mathbb{P}_P^2 \subset \overline{Q}$, parametrizing lines through a point $P \in \mathbb{P}(\mathbb{C}^4)$, and a $\mathbb{P}_V^2 \in \overline{Q}$, parametrizing the lines in a plane $V \cong \mathbb{P}^2 \subset \mathbb{P}(\mathbb{C}^4)$. Another basis is given by \mathbb{P}_V^2 and H^2 , with H the hyperplane section.

The (strict) pull back of \mathbb{P}_V^2 along P is $V \times V \subset \mathbb{P}^3 \times \mathbb{P}^3$ and this pulls back to $Z := (2\Theta_Y) \times (2\Theta_Y)$ in $Y \times Y$. The pull-back of H is $2\Theta_X = Y \times (2\Theta_Y) + (2\Theta_Y) \times Y$, twice the product polarization on $Y \times Y$. For general $\tau \in \mathbb{H}_4$, one has $D^2(X_\tau) = \langle \Theta_\tau^2 \rangle$, which specializes to $\langle \Theta_X^2 \rangle$. Therefore we must show that

$$[Z] \notin \langle \Theta_X^2, [\Delta], [\Delta^-] \rangle \subset H^4(Y \times Y, \mathbb{Q}) = \bigoplus_a H^a(Y, \mathbb{Q}) \otimes H^{4-a}(Y, \mathbb{Q}).$$

Note that $[Z] \in H^2 \otimes H^2$ (with $H^k := H^k(Y, \mathbb{Q})$). On the other hand,

$$\langle \Theta_X^2, [\Delta], [\Delta^-] \rangle \cap H^2 \otimes H^2 = \langle \Theta_X^2 - ([\Delta] + [\Delta^-]) \rangle,$$

in fact, $\Theta_X^2 = 2(\{P\} \times Y + \Theta_Y \times \Theta_Y + Y \times \{P\})$ has no $(3, 1)$, $(1, 3)$ Künneth components, and $[\Delta] + [\Delta^-] = 2(\{P\} \times Y + \delta_{2,2} + Y \times \{P\})$, with $[\Delta] = \sum \delta_{a,b}$ the Künneth decomposition of $[\Delta]$. Since $\delta_{2,2}$ has a non-trivial component in $H^{2,0}(Y) \otimes H^{0,2}(Y)$, whereas $[Z] = [\Theta_Y \times \Theta_Y] \in H^{1,1} \otimes H^{1,1}$ has a trivial component there, we conclude that $[Z] \notin \langle \Theta_X^2, [\Delta], [\Delta^-] \rangle$.

By specialization, this proves that $[Z] \notin \langle \Theta_{X_\tau}^2 \rangle$ for any X_τ . \square

4 An isogeny

4.1 In this section we give another description of the map K_- from theorem 3.7, using an isogeny on X_τ . This new description relates the quadratic relation between certain theta functions to geometrical properties of X_τ , see 4.4.

4.2 A basis of $H^0(X, \mathcal{O}(4\Theta))$ is given by the $4^g = 2^{2g}$ functions

$$\theta \begin{bmatrix} \varepsilon \\ \varepsilon' \end{bmatrix}(\tau, 2z) \quad \varepsilon, \varepsilon' \in \{0, 1\}^g.$$

Let $V \subset H^0(X, \mathcal{O}(4\Theta))$ be the subspace spanned by the six even theta functions with vanishing Nullwert:

$$V := \langle \dots, \theta \begin{bmatrix} \sigma & \sigma \\ \sigma' & \sigma' \end{bmatrix}(\tau, 2z), \dots \rangle_{(\sigma, \sigma' \in \{0, 1\}^2, \sigma \sigma' = 1)} \subset H^0(X, \mathcal{O}(4\Theta)).$$

4.3 Proposition. *The image of $X = X_\tau$ under the rational map:*

$$\Phi_V : X \rightarrow \mathbb{P}V, \quad z \mapsto (\dots : \theta \begin{bmatrix} \sigma & \sigma \\ \sigma' & \sigma' \end{bmatrix}(\tau, 2z) : \dots)$$

is a smooth quadric $\overline{Q} \subset \mathbb{P}^5$. The quadratic relation on the six theta functions is:

$$\left(\theta^2 \begin{bmatrix} 1010 \\ 1010 \end{bmatrix} - \theta^2 \begin{bmatrix} 1010 \\ 1111 \end{bmatrix} - \theta^2 \begin{bmatrix} 1111 \\ 1010 \end{bmatrix} + \theta^2 \begin{bmatrix} 1111 \\ 0101 \end{bmatrix} - \theta^2 \begin{bmatrix} 0101 \\ 0101 \end{bmatrix} + \theta^2 \begin{bmatrix} 0101 \\ 1111 \end{bmatrix} \right) (\tau, 2z) = 0.$$

There is a commutative diagram:

$$\begin{array}{ccc} X & \xrightarrow{K_-} & \overline{Q} \subset \mathbb{P}^5 \\ \uparrow 1-\phi & \nearrow \Phi_V & \\ X & & \end{array}$$

Proof. We consider the subspace $\Gamma(\mathcal{O}(2\Theta))_-$ where $\phi^* = -I$. Remark 2.5 shows that the functions $\theta \begin{bmatrix} \varepsilon & \varepsilon \\ \varepsilon' & \varepsilon' \end{bmatrix}(\tau, z(I-A))$ give a basis of $\Gamma(\mathcal{O}(2\Theta))_-$. From formula 2.3.1 we have:

$$\theta \begin{bmatrix} \varepsilon & \varepsilon \\ \varepsilon' & \varepsilon' \end{bmatrix}(\tau, zA) = (-1)^{\varepsilon' \varepsilon'} \theta \begin{bmatrix} \varepsilon & \varepsilon \\ \varepsilon' & \varepsilon' \end{bmatrix}(\tau, z),$$

and thus a basis for $\Gamma(\mathcal{O}(2\Theta))_-$ is given by the $2^{n-1}(2^n - 1)$ functions $\theta \begin{bmatrix} \varepsilon & \varepsilon \\ \varepsilon' & \varepsilon' \end{bmatrix}(\tau, z(I-A))$ with $\varepsilon' \varepsilon' \equiv 1 \pmod{2}$. Since $(I-A)^2 = I - 2A + A^2 = -2A$ we get:

$$V = (1 - \phi)^* \Gamma(2\Theta)_-.$$

In the case $n = 2$ we used the basis with the six Y_{ab} 's. One easily finds, with $\Theta[\sigma] := \theta[\begin{smallmatrix} \sigma \\ 0 \end{smallmatrix}](2\tau, 2z)$:

$$\begin{aligned} \theta \begin{bmatrix} 1010 \\ 1010 \end{bmatrix} (\tau, z(I-A)) &= -\Theta[0010] - \Theta[0111] + \Theta[1000] + \Theta[1101] \\ &= -Y_{02} - Y_{13}, \\ \theta \begin{bmatrix} 1010 \\ 1111 \end{bmatrix} (\tau, z(I-A)) &= -\Theta[0010] + \Theta[0111] + \Theta[1000] - \Theta[1101] \\ &= -Y_{02} + Y_{13}, \end{aligned}$$

similarly one has:

$$\begin{aligned} \theta \begin{bmatrix} 0101 \\ 0101 \end{bmatrix} (\tau, z(I-A)) &= -Y_{01} - Y_{23} & \theta \begin{bmatrix} 1111 \\ 1010 \end{bmatrix} (\tau, z(I-A)) &= -Y_{03} + Y_{12} \\ \theta \begin{bmatrix} 0101 \\ 1111 \end{bmatrix} (\tau, z(I-A)) &= -Y_{01} + Y_{23} & \theta \begin{bmatrix} 1111 \\ 0101 \end{bmatrix} (\tau, z(I-A)) &= -Y_{03} - Y_{12}. \end{aligned}$$

From these equations one can express the Y_{ab} 's in the $\theta[\begin{smallmatrix} \epsilon & \epsilon \\ \epsilon' & \epsilon' \end{smallmatrix}](\tau, z(I-A))$. The quadratic relation between the Y_{ab} 's then gives the relation in the proposition. \square

4.4 Remark. The existence of a quadratic relation between the six even theta functions with a vanishing Nullwert can also be obtained from the following observations.

First of all, each vanishing Nullwert corresponds to a singular point on the theta divisor Θ of X , the singular point is a point of order two in X . Thus Θ has at least six singular points. Furthermore, for $x \in \text{Sing}(\Theta)$ we have that

$$f_x := \theta_{0,0}(\tau, z-x)\theta_{0,0}(\tau, z+x) \in \Gamma_{00} := \{s \in H^0(X, \mathcal{O}(2\Theta)) : \mu_0(s) \geq 4\},$$

where $\mu_0(s)$ is the multiplicity at $0 \in X$ of the section s . The six functions $f_x \in \Gamma_{00}$ we obtain are the six theta functions $\theta[\begin{smallmatrix} \sigma & \sigma \\ \sigma' & \sigma' \end{smallmatrix}]^2(\tau, z)$ with $\sigma'\sigma = 1$.

Since the general X we consider is not a product of lower dimensional ppav's, one has (see [vGvdG]):

$$\dim \Gamma_{00} = 2^4 - 1 - 10 = 5.$$

Thus the six functions $\theta[\begin{smallmatrix} \sigma & \sigma \\ \sigma' & \sigma' \end{smallmatrix}]^2(\tau, z) \in \Gamma_{00}$ are linearly dependent and therefore one has also a linear relation between the six $\theta[\begin{smallmatrix} \sigma & \sigma \\ \sigma' & \sigma' \end{smallmatrix}]^2(\tau, 2z)$.

The results of Debarre [D] imply that $\text{Sing}(\Theta_\tau)$, for general $\tau \in \mathbb{H}_4$ consists of exactly 6 points (that the general X_τ is not the jacobian of a hyperelliptic curve follows from the fact that the sum of any three vanishing theta characteristics of X_τ is even, rather than odd).

Geometrically, the argument above shows that $K(\text{Sing}(\Theta))$, which in general consists of 6 points, spans at most a $\mathbb{P}^4 \subset \mathbb{P}^{15}$, in fact the map K may also be given by (cf. [vGvdG]):

$$K : X_\tau \rightarrow \mathbb{P}H^0(X_\tau, \mathcal{O}(2\Theta_\tau))^*, \quad x \mapsto \theta_{0,0}(\tau, z-x)\theta_{0,0}(\tau, z+x).$$

4.5 Remark. We indicate an alternative derivation of the quadratic relation in the previous proposition, using the identities (with $\sigma_1, \sigma_2 \in \{0, 1\}^2$):

$$\theta \begin{bmatrix} \sigma_1 & \sigma_2 \\ 0 & 0 \end{bmatrix} (2\tau, 0) = \theta \begin{bmatrix} \sigma_2 & \sigma_1 \\ 0 & 0 \end{bmatrix} (2\tau, 0) \quad \tau \in \mathbb{H}_{2n}.$$

The multiplication formula for theta functions ([I], Chapter IV, Thm. 2, p.139) shows that the expression for $\theta^2 \begin{bmatrix} \varepsilon & \varepsilon \\ \varepsilon' & \varepsilon' \end{bmatrix} (\tau, z)$ in terms of the standard basis of $\Gamma(\mathcal{O}(2\Theta_\tau))$ is given by:

$$\begin{aligned} & \theta \begin{bmatrix} \varepsilon & \varepsilon \\ \varepsilon' & \varepsilon' \end{bmatrix}^2 (\tau, z) \\ &= \sum_{\sigma_1, \sigma_2 \in \{0, 1\}^2} (-1)^{(\sigma_1, \sigma_2)' \varepsilon'} \theta \begin{bmatrix} \sigma_1 + \varepsilon & \sigma_2 + \varepsilon \\ 0 & 0 \end{bmatrix} (2\tau, 0) \theta \begin{bmatrix} \sigma_1 & \sigma_2 \\ 0 & 0 \end{bmatrix} (2\tau, 2z). \end{aligned}$$

It is now a straight-forward computation to verify the relation. One can also use the proof of 3.6 as follows. Define:

$$B_{Q_m}(X, Y) := \sum_{\sigma} (-1)^{t_{\sigma \varepsilon'}} X_{\sigma} Y_{\sigma + \varepsilon}, \quad \text{with } m = \begin{bmatrix} \varepsilon \\ \varepsilon' \end{bmatrix}$$

an even theta characteristic. Note that B_{Q_m} is just the bilinear form associated to the quadratic form Q_m in the proof of Proposition 3.6. Upon substituting $Y_{\sigma} := \theta \begin{bmatrix} \sigma \\ 0 \end{bmatrix} (2\tau, 0)$ one obtains linear forms in the X_{σ} , these 6 forms are linearly dependent exactly when there is a linear relation between the $\theta^2 \begin{bmatrix} \varepsilon & \varepsilon \\ \varepsilon' & \varepsilon' \end{bmatrix} (\tau, z)$.

In the proof of 3.6, we showed that a certain linear combination Q of the six Q_m 's is a quadric in the coordinates Y_{st} of \mathbb{P}^5 . Thus Q is a linear combination of terms $(X_{st} - X_{ts})(X_{uv} - X_{vu})$, and therefore its associated bilinear form B_Q is a linear combination of terms $(Y_{uv} - Y_{vu})X_{st}$. Since we have $\theta \begin{bmatrix} \sigma_1 & \sigma_2 \\ 0 & 0 \end{bmatrix} = \theta \begin{bmatrix} \sigma_2 & \sigma_1 \\ 0 & 0 \end{bmatrix}$, upon substituting $Y_{\sigma} := \theta \begin{bmatrix} \sigma \\ 0 \end{bmatrix} (2\tau, 0)$ in B_Q we get zero.

5 Prym varieties

5.1 Let C_{n+1} be a smooth curve of genus $n+1$. A subgroup $G \cong \mathbb{Z}/4\mathbb{Z} \subset \text{Pic}^0(C_{n+1})$ defines a cyclic étale 4:1 covering of C_{n+1} and an intermediate étale 2:1 cover:

$$C_{n+1} \leftarrow C_{2n+1} \leftarrow C_{4n+1}.$$

The Prym variety P of the C_{4n+1}/C_{2n+1} is a principally polarized abelian variety of dimension $2n$. The covering automorphism of C_{4n+1} over C_{n+1} induces an action of $\mathbb{Z}[i]$ on P .

5.2 Schoen constructs cycles on P by taking the inverse image of $\mathbb{P}^n = |K| \subset C_{n+1}^{(2n)}$ in $C_{4n+1}^{(2n)}$ (with $C^{(i)} = \text{Sym}^i C$), which is reducible, and mapping the components to P . He shows that suitable linear combinations of these cycles span the space of Weil-Hodge cycles $W \subset H^{2n}(P, \mathbb{Q})$.

In the case $n = 2$ we show below that the general $X_{\tau}, \tau \in \mathbb{H}_4$, is such a Prym variety.

5.3 Theorem. *Let P be as in 5.1. Then:*

1. *The abelian variety P is of Weil type (n, n) , with field $\mathbb{Q}(i)$ and $\det H = 1$.*
2. *There exist bases of $H_1(P, \mathbb{Z})$ and $H_1(P, \mathbb{R})$ (which has the structure of a complex vector space via its identification with T_0P) such that the period matrix Ω satisfies:*

$$\Omega = \begin{pmatrix} I \\ \tau \end{pmatrix} : H_1(P, \mathbb{Z}) \cong \mathbb{Z}^{4n} \rightarrow H_1(X, \mathbb{R}) = T_0P \cong \mathbb{C}^{2n} \quad \text{with } \tau \in \mathbb{H}_{2n}.$$

In particular, P is isomorphic to an X_τ with $\tau \in \mathbb{H}_{2n}$.

3. *In case $n = 2$, the general abelian variety of Weil type $(2, 2)$ and field $\mathbb{Q}(i)$ with $\det H = 1$ is isogeneous to a Prym variety as in 5.1.*

Proof. We consider the action of the automorphism ϕ of order 4 on $H_1(P, \mathbb{Z})$. According to [F], p. 62, there is a symplectic basis \hat{A}_i, \hat{B}_i ($0 \leq i \leq 4n$) of $H_1(C_{4n+1}, \mathbb{Z})$ with:

$$\phi^k(\hat{A}_i) = \hat{A}_{i+kn}, \quad \phi^k(\hat{B}_i) = \hat{B}_{i+kn} \quad (1 \leq i \leq n).$$

Since $H_1(P, \mathbb{Z}) = H_1(C_{4n+1}, \mathbb{Z})^{\phi_* = -I}$ (the classes anti-invariant under ϕ_*), a basis of $H_1(P, \mathbb{Z})$ is given by

$$\alpha_i := \hat{A}_i - \hat{A}_{i+2n}, \quad \beta_i := \hat{B}_i - \hat{B}_{i+2n} \quad (1 \leq i \leq 2n).$$

An easy computation shows that the action of ϕ on $H_1(P, \mathbb{Z})$ is given by the matrix M , i.e. $\phi(\alpha_i) = \alpha_{i+n}$, $\phi(\beta_i) = \beta_{i+n}$ with $1 \leq i \leq n$. Taking the α_i as a \mathbb{C} -basis of $H_1(P, \mathbb{R})$, the period matrix must satisfy $M\Omega = \Omega A$, which implies that $\tau \in \mathbb{H}_{2n}$. This proves the second point.

Using this result, the first point is easy to verify (cf. 2.1).

For the last point, we consider the Prym map:

$$\mathcal{P} : \mathcal{M}_{3, \mathbb{Z}/4\mathbb{Z}} \rightarrow \mathcal{A}_4, \quad (C, G) \mapsto P$$

where $\mathcal{M}_{3, \mathbb{Z}/4\mathbb{Z}}$ is the moduli space of genus 3 curves with a cyclic subgroup of order 4 in Pic^0 and \mathcal{A}_4 is the moduli space of principally polarized abelian varieties. We already proved that $\text{Im}(\mathcal{P})$ lies in the 4-fold $\mathcal{H} := \text{Im}(\mathbb{H}_4 \rightarrow \mathbb{S}_4 \rightarrow \mathcal{A}_4)$. To show that $\text{Im}(\mathcal{P})$ is Zariski dense in \mathcal{P} it suffices to show that the differential of \mathcal{P} has rank 4 at some point of $\mathcal{M}_{3, \mathbb{Z}/4\mathbb{Z}}$, or, equivalently, that the codifferential of \mathcal{P} is injective at some point of \mathcal{H} .

The remainder follows the arguments of Schoen, [S], p. 26–p. 30, we just show the (obvious) modifications. The cotangent space of $\mathcal{M}_{3, \mathbb{Z}/4\mathbb{Z}}$ at a smooth point $[(C, G)]$ is $H^0(C, \Omega_C^{1 \otimes 2})$. Let $\pi : C_9 \rightarrow C$ be the 4:1 étale map, then, using the action of ϕ , we have:

$$\pi_* \Omega_{C_9}^1 = \Omega_C^1 \oplus (\Omega_C^1 \otimes \alpha^{\otimes 2}) \oplus (\Omega_C^1 \otimes \alpha) \oplus (\Omega_C^1 \otimes \alpha^{\otimes 3}) \quad \text{with } G = \langle \alpha \rangle.$$

Then $T_P^* \mathcal{A}_4$ and $T_P^* \mathcal{H}$, the cotangent spaces of \mathcal{A}_4 and \mathcal{H} at P are given by:

$$\begin{aligned} T_P^* \mathcal{A}_4 &= \text{Sym}^2(H^0(C, \Omega_C^1 \otimes \alpha) \oplus H^0(C, \Omega_C^1 \otimes \alpha^3)), \\ T_P^* \mathcal{H} &= H^0(C, \Omega_C^1 \otimes \alpha) \otimes H^0(C, \Omega_C^1 \otimes \alpha^3). \end{aligned}$$

The codifferential of $\mathcal{P} : \mathcal{M}_3, \mathbb{Z}/4\mathbb{Z} \rightarrow \mathcal{H}$ is now just the multiplication map:

$$H^0(C, \Omega_C^1 \otimes \alpha) \otimes H^0(C, \Omega_C^1 \otimes \alpha^3) \rightarrow H^0(C, \Omega_C^{1 \otimes 2}).$$

The proof that this map is injective for general $[C] \in \mathcal{M}_3$ follows from the Base Point Free Pencil trick, as in [S], p. 30. \square

5.4 Remark. Since $\dim \mathcal{M}_4 = 9 = 3^2$, one might hope that the Prym map: $\mathcal{P} : \mathcal{M}_4, \mathbb{Z}/4\mathbb{Z} \rightarrow \mathcal{A}_6$ would be generically finite on its image. If that is the case, then $\text{Im}(\mathcal{P})$ is Zariski dense in \mathcal{H}_6 and Schoen's cycles would solve the Hodge conjecture for the general abelian sixfold of Weil-type with field $\mathbb{Q}(i)$ and $\det H = 1$.

5.5 Remark. The previous Proposition also 'explains' the 6 singular points of order two on the theta divisor of the abelian fourfolds we considered. Recall that a theta characteristic on a curve C is a line bundle L with $L^{\otimes 2} = \Omega_C^1$.

For a curve of genus $n+1$ and a point $\beta \in \text{Pic}^0(C)$ of order two, there are $2^{n-1}(2^n-1)$ pairs of odd theta characteristics L, L' with $L' = L \otimes \beta$ (this is most easily checked using the classical notation for characteristics). Let $\pi' : C' \rightarrow C$ be the étale 2:1 cover defined by β , then

$$\pi'^* L \cong \pi'^* L', \quad H^0(C', \pi'^* L) = H^0(C, L) \oplus H^0(C, L'), \quad (\pi'^* L)^{\otimes 2} = \Omega_C^1,$$

thus $\pi'^* L$ is an even theta characteristic with at least two independent sections.

Let now $\alpha \in \text{Pic}^0(C)$ with $\alpha^{\otimes 2} = \beta$. Then $\gamma := \pi'^* \alpha$ has order two in $\text{Pic}(C')$ and moreover, by Serre duality, $H^0(C, L \otimes \alpha) \cong H^0(C, L \otimes \alpha^{-1})^* = H^0(C, L' \otimes \alpha)^*$. Therefore, also $(\pi'^* L) \otimes \gamma = \pi'^*(L \otimes \alpha)$ is an even theta characteristic on C' .

Now it is well known (for example from the classical Schottky–Jung identities) that pairs $M, M \otimes \gamma$ of even theta characteristics on C' with $H^0(C', M) > 0$ correspond to singular points, of order two and even multiplicity, on the theta divisor of the Prym variety of the covering $C'' \rightarrow C'$ defined by γ . These singular points correspond precisely to the vanishing (even) theta nulls.

Starting from a genus $n+1$ curve, the $2n$ -dimensional Prym varieties introduced in 5.1, thus have at least $2^{n-1}(2^n-1)$ vanishing (even) theta nulls, in agreement with the fact that X_τ , for $\tau \in \mathbb{H}_{2n}$ has that number of vanishing theta nulls (simply put $z = 0$ in 2.3.1).

References

- [D] Debarre, O.: Annulation de thétaconstantes sur les variétés abéliennes de dimension quatre. C. R. Acad. Sci. Paris, t. **305**, Série I, 885–888 (1987)
- [F] Fay, J.D.: Theta Functions on Riemann Surfaces LNM. **352**, Springer-Verlag (1973)

- [vG1] van Geemen, B.: Projective models of Picard modular varieties. In: Ballico, E. et al., (Eds) *Classification of Irregular Varieties, proceedings Trento 1990*. Springer LNM **1515**, (1992)
- [vG2] van Geemen, B.: An introduction to the Hodge conjecture for abelian varieties. preprint Utrecht University (1993)
- [vGvdG] van Geemen, B., van der Geer, G.: Kummer varieties and the moduli spaces of abelian varieties. *Am. J. of Math.* **108**, 615–642 (1986)
- [I] Igusa, J.I.: *Theta Functions*. Springer-Verlag, Die Grundlehren der mathematischen Wissenschaften. Band 194, (1972)
- [S] Schoen, C.: Hodge classes on self-products of a variety with an automorphism. *Comp. Math.* **65**, 3–32 (1988)
- [W] Weil, A.: Abelian varieties and the Hodge ring. In: *Collected Papers, Vol. III*, 421–429. Springer-Verlag (1980)

